

Three types of generalized Kadomtsev–Petviashvili equations arising from baroclinic potential vorticity equation*

Zhang Huan-Ping(张焕萍)^{a)}, Li Biao(李彪)^{a)d)†},
Chen Yong(陈勇)^{a)b)}, and Huang Fei(黄菲)^{c)}

^{a)}Nonlinear Science Center and Department of Mathematics, Ningbo University, Ningbo 315211, China

^{b)}Institute of Theoretical Computing, East China Normal University, Shanghai 200062, China

^{c)}Department of Marine Meteorology, Ocean University of China, Qingdao 266003, China

^{d)}Key Laboratory of Mathematics Mechanization, Chinese Academy of Sciences, Beijing 100190, China

(Received 24 March 2009; revised manuscript received 24 July 2009)

By means of the reductive perturbation method, three types of generalized (2+1)-dimensional Kadomtsev–Petviashvili (KP) equations are derived from the baroclinic potential vorticity (BPV) equation, including the modified KP (mKP) equation, standard KP equation and cylindrical KP (cKP) equation. Then some solutions of generalized cKP and KP equations with certain conditions are given directly and a relationship between the generalized mKP equation and the mKP equation is established by the symmetry group direct method proposed by Lou *et al.* From the relationship and the solutions of the mKP equation, some solutions of the generalized mKP equation can be obtained. Furthermore, some approximate solutions of the baroclinic potential vorticity equation are derived from three types of generalized KP equations.

Keywords: baroclinic potential vorticity equation, generalized Kadomtsev–Petviashvili equation, symmetry groups, approximate solution

PACC: 0220, 1130, 3310c, 0340k

1. Introduction

In recent years, the study of nonlinear partial differential equations (PDEs) have become one of the most exciting and extremely active areas of research. Many methods can be used to obtain solutions of PDEs, such as symmetry reductions, the general direct method and the extended Jacobi elliptic function rational expansion method, and so on.^[1–10] However, there are many difficulties in obtaining solutions of baroclinic potential vorticity (BPV) equation^[11] by these methods. Luckily, the reductive perturbation method can make an intricate equation become another new equation, then the intricate equation can be researched more successfully with the help of a new equation. This method has been used in many fields. Lou *et al.*^[12] derived coupled KdV equations from two-layer fluids. Tang *et al.*^[13] obtained the variable coefficient KdV equation from the Euler equation with an earth rotation term. According to the KdV equation,

they obtained the approximate solution of the Euler equation with an earth rotation term. Gao *et al.*^[14] presented a coupled variable coefficient modified KdV equation from a two-layer fluid system. The main purpose of this paper is to obtain the approximate solutions of the BPV equation. The paper is organized as follows. In Section 2, we obtain generalized modified Kadomtsev–Petviashvili (mKP), KP and cylindrical KP (cKP) equations from the BPV equation by means of the reductive perturbation method. Some KP equations have been studied by some authors. Li *et al.*^[15] obtained soliton-like solution and periodic form solution of a KP equation. Liu^[16] gained a solution of the cKP equation by auto-Backlund transformation. Wazwaz^[17] obtained multiple-soliton solutions of the mKP equation by the Hirota’s bilinear method. In Section 3, under certain conditions we obtain the solutions of the generalized cKP and KP equation directly, and then get an approximate solution of the BPV equation. In Section 4, by making use of the

*Project supported by National Natural Science Foundation of China (Grant Nos. 10735030 and 40775042), Ningbo Natural Science Foundation (Grant No. 2008A610017), National Basic Research Program of China (973 Program) (Grant Nos. 2005CB422301 and 2007CB814800) and K.C. Wong Magna Fund in Ningbo University.

†Corresponding author. E-mail: biaolee2000@yahoo.com.cn

© 2010 Chinese Physical Society and IOP Publishing Ltd

<http://www.iop.org/journals/cpb> <http://cpb.iphy.ac.cn>

symmetry group direct method proposed by Lou *et al.*,^[18–21] we establish a relationship between the generalized mKP equation and mKP equation, and then based on this relationship we obtain an approximate solution of the BPV equation by complex and tedious calculations. In the last section, we give the conclusion of the paper.

2. Generalized mKP, KP and cKP equations from (3+1)-dimensional BPV equation

The (3 + 1)-dimensional BPV equation may be written as

$$q_t + \psi_x q_y - \psi_y q_x + \beta \psi_x = 0, \quad (1)$$

$$q = \psi_{xx} + \psi_{yy} + \psi_{zz}, \quad (2)$$

where β is constant, $\psi \equiv \psi(x, y, z, t)$, $q \equiv q(x, y, z, t)$, and subscripts x, y, z, t represent partial derivatives.

First, by substituting Eq. (2) into Eq. (1), BVP equation becomes:

$$\begin{aligned} &\psi_{txx} + \psi_{tyy} + \psi_{tzz} + \psi_x \psi_{xxy} + \psi_x \psi_{yyx} + \psi_x \psi_{yzz} \\ &- \psi_y \psi_{xx} - \psi_y \psi_{xyy} - \psi_y \psi_{xzz} + \beta \psi_x = 0. \end{aligned} \quad (3)$$

Now we use the long wave approximation in the x -direction and z -direction and assume the stream functions ψ to be

$$\psi = \psi_0(y, t) + \psi_1(x, y, z, t). \quad (4)$$

We introduce the stretched variables

$$X = \epsilon(x - c_0 t), \quad Z = \epsilon^2(z - c_1 t), \quad T = \epsilon^3 t, \quad (5)$$

where $X \equiv X(x, t)$, $Z \equiv Z(z, t)$, $T \equiv T(t)$, c_0 and c_1 are constants, ϵ is a small parameter. In general, the base field $\psi_0(y, t)$ is often taken only as a linear function of y , $\psi_1(x, y, z, t)$ is expanded as

$$\psi_1(x, y, z, t) = \sum_{i=1}^{\infty} \epsilon^i \phi_i(X, y, Z, T) \equiv \sum_{i=1}^{\infty} \epsilon^i \phi_i. \quad (6)$$

We also have the expansion

$$\psi_0(y, t) = U_0(y) + \sum_{i=1}^{\infty} \epsilon^i U_i(y, T). \quad (7)$$

Then substituting Eqs. (5), (6) and (7) into Eq. (3), we obtain

$$M_1 \epsilon^2 + M_2 \epsilon^3 + M_3 \epsilon^4 + O(\epsilon^5) = 0, \quad (8)$$

where

$$\begin{aligned} M_1 = & -U_{0y}(y)\phi_{1Xyy} + U_{0yyy}(y)\phi_{1X} \\ & - c_0\phi_{1Xyy} + \beta\phi_{1X}, \end{aligned}$$

$$\begin{aligned} M_2 = & \phi_{1X}\phi_{1yyy} - c_0\phi_{2Xyy} - U_{1y}(y, T)\phi_{1Xyy} \\ & - \phi_{1Xyy}\phi_{1y} - c_1\phi_{1yYZ} + U_{1yyy}(y, T)\phi_{1X} \\ & - U_{0y}(y)\phi_{2Xyy} + U_{0yyy}(y)\phi_{2X} + \beta\phi_{2X}, \end{aligned}$$

$$\begin{aligned} M_3 = & \phi_{2X}\phi_{1yyy} - c_0\phi_{3Xyy} + U_{1yyy}(y, T)\phi_{2X} \\ & + \beta\phi_{3X} + \phi_{1Tyy} - U_{0y}(y)\phi_{1XXX} \\ & + U_{0yyy}(y)\phi_{3X} - U_{2y}(y, T)\phi_{1Xyy} - c_0\phi_{1XXX} \\ & - U_{0y}(y)\phi_{3Xyy} - U_{1y}(y, T)\phi_{2Xyy} - c_1\phi_{2yYZ} \\ & + U_{2yyy}\phi_{1X} + \phi_{1X}\phi_{2yyy} + U_{1yyT}(y, T) \\ & - \phi_{2Xyy}\phi_{1y} - \phi_{1Xyy}\phi_{2y}. \end{aligned}$$

Vanishing the ϵ^2 of Eq. (8) by Maple, we get a special solution

$$\phi_1 = A(X, Z, T)G(y, T) \equiv AG, \quad (9)$$

where G is determined by

$$U_{0yyy}(y)G - U_{0y}(y)G_{yy} - c_0G_{yy} + \beta G = 0. \quad (10)$$

Then vanishing the ϵ^3 of Eq. (8), we have

$$\begin{aligned} \phi_2 = & G_1(y, T)A(X, Z, T)^2 + G_2(y, T)A(X, Z, T) \\ & + G_3(y, T) \int A_Z(X, Z, T) dX \\ \equiv & G_1 A^2 + G_2 A + G_3 \int A_Z dX, \end{aligned} \quad (11)$$

where G_1, G_2 and G_3 should satisfy

$$\begin{aligned} &-G_{yy}G_y + GG_{yyy} + 2U_{0yyy}(y)G_1 - 2U_{0y}(y)G_{1yy} \\ &- 2c_0G_{1yy} + 2\beta G_1 = 0, \\ &U_{1yyy}(y, T)G - U_{1y}(y, T)G_{yy} + \beta G_2 + U_{0yyy}(y)G_2 \\ &- U_{0y}(y)G_{2yy} - c_0G_{2yy} = 0, \\ &U_{0yyy}(y)G_3 - U_{0y}(y)G_{3yy} - c_0G_{3yy} - c_1G_{yy} \\ &+ \beta G_3 = 0. \end{aligned} \quad (12)$$

Substituting the solutions of Eqs. (10), (12) and $\phi_3 = 0$ into M_3 , integrating the result with respect to y from 0 to y_0 , then we obtain

$$\begin{aligned} &a_1 A_T + a_2 A_{XXX} + a_3 A A_X + a_4 A A_Z + a_5 A^2 A_X \\ &+ a_6 A_X \int A_Z dX + a_7 \int A_{ZZ} dX \\ &+ a_8 A_X + a_9 A_Z + a_{10} A + a_{11} = 0, \end{aligned} \quad (13)$$

where $a_i \equiv a_i(T)$, $i = 1, 2, \dots, 11$

$$a_1 = \int_0^{y_0} \int_0^{y_1} G_{yy} dy dy_1,$$

$$\begin{aligned}
 a_2 &= \int_0^{y_0} \int_0^{y_1} (-U_{0y}(y)G - c_0G) dy dy_1, \\
 a_3 &= \int_0^{y_0} \int_0^{y_1} [2U_{1yyy}(y, T)G_1 - 2U_{1y}(y, T)G_{1yy} \\
 &\quad - G_y G_{2yy} + G_{yyy}G_2 + GG_{2yyy} \\
 &\quad - G_{yy}G_{2y}] dy dy_1, \\
 a_4 &= \int_0^{y_0} \int_0^{y_1} (G_{yyy}G_3 - 2c_1G_{1yy} - G_y G_{3yy}) dy dy_1, \\
 a_5 &= \int_0^{y_0} \int_0^{y_1} (-2G_y G_{1yy} + 2G_{yyy}G_1 + GG_{1yyy} \\
 &\quad - G_{yy}G_{1y}) dy dy_1, \\
 a_6 &= \int_0^{y_0} \int_0^{y_1} (GG_{3yyy} - G_{yy}G_{3y}) dy dy_1, \\
 a_7 &= \int_0^{y_0} \int_0^{y_1} (-c_1G_{3yy}) dy dy_1, \\
 a_8 &= \int_0^{y_0} \int_0^{y_1} [-U_{2y}(y, T)G_{yy} + U_{2yyy}(y, T)G \\
 &\quad - U_{1y}(y, T)G_{2yy} + U_{1yyy}(y, T)G_2] dy dy_1, \\
 a_9 &= \int_0^{y_0} \int_0^{y_1} [U_{1yyy}(y, T)G_3 - c_1G_{2yy} \\
 &\quad - U_{1y}(y, T)G_{3yy}] dy dy_1, \\
 a_{10} &= \int_0^{y_0} \int_0^{y_1} G_{yyT} dy dy_1, \\
 a_{11} &= \int_0^{y_0} \int_0^{y_1} U_{1yyT}(y, T) dy dy_1.
 \end{aligned}$$

Equation (13) contains the following important cases:

(i) If $a_4 = 0, a_5 = 0, a_6 = 0, a_8 = 0, a_9 = 0, a_{10} = 0, a_{11} = 0$, equation (13) becomes a generalized KP equation

$$\begin{aligned}
 a_1 A_T + a_2 A_{XXX} + a_3 AA_X \\
 + a_7 \int A_{ZZ} dX = 0. \quad (14)
 \end{aligned}$$

(ii) If $a_4 = 0, a_5 = 0, a_6 = 0, a_8 = 0, a_9 = 0, a_{10} = 0$, equation (13) becomes a generalized cKP equation

$$\begin{aligned}
 a_1 A_T + a_2 A_{XXX} + a_3 AA_X \\
 + a_7 \int A_{ZZ} dX + a_{11} A = 0. \quad (15)
 \end{aligned}$$

(iii) If $a_3 = 0, a_4 = 0, a_8 = 0, a_9 = 0, a_{10} = 0, a_{11} = 0$, equation (13) becomes a generalized mKP equation

$$\begin{aligned}
 a_1 A_T + a_2 A_{XXX} + a_5 A^2 A_X \\
 + a_6 A_X \int A_Z dX + a_7 \int A_{ZZ} dX = 0. \quad (16)
 \end{aligned}$$

In the next section, we discuss the solutions of Eq. (3) with the help of Eqs. (14), (15) and (16) by a direct method and symmetry group direct method.

3. The solutions of BPV equation from generalized KP and cKP equations

In this section, we divide into two parts to obtain approximate solutions of the BPV equation with the help of the generalized KP and cKP equations.

First, we set about the generalized KP equation to obtain the approximate solution of Eq. (3). We know that equation (14) should satisfy

$$\begin{aligned}
 a_4 = a_5 = a_6 = a_8 = a_9 = a_{10} = a_{11} = 0, \\
 U_{0yyy}(y)A_X G - U_{0y}(y)A_X G_{yy} - c_0 A_X G_{yy} = 0, \\
 -G_{yy}G_y + GG_{yyy} + 2U_{0yyy}(y)G_1 - 2U_{0y}(y)G_{1yy} \\
 - 2c_0 G_{1yy} = 0, \\
 U_{0yyy}(y)G_3 - U_{0y}(y)G_{3yy} - c_0 G_{3yy} - c_1 G_{yy} = 0, \\
 U_{1yyy}(y, T)G - U_{1y}(y, T)G_{yy} + \beta_1 G \\
 + U_{0yyy}(y)G_2 - U_{0y}(y)G_{2yy} - c_0 G_{2yy} = 0. \quad (17)
 \end{aligned}$$

According to Eq. (17), we obtain $G(y, k), G_1(y, k), G_2(y, k), G_3(y, k), U_0(y), U_1(y, k)$ and $U_2(y, k)$ after complicated computing process

$$\begin{aligned}
 G = C_8, \quad G_1 = C_7, \quad G_2 = F_3(T)(C_1 y + C_2), \\
 G_3 = c_1 F_3(T) + F_4(T)(C_1 y + C_2), \\
 U_0(y) = -\frac{1}{6}\beta y^3 + \frac{1}{2}C_4 y^2 + C_5 y + C_6, \\
 U_i(y, k) = 0 \quad (i \geq 2), \\
 U_1(y, T) = \frac{1}{2}C_1 y^2 + C_2 y + C_3 + F_1(T), \quad (18)
 \end{aligned}$$

where C_i ($i = 1, 2, \dots, 8$) are arbitrary constants, $F_1(T), F_3(T)$ and $F_4(T)$ are arbitrary functions with respect to time. At the same time, we can obtain a_1, a_2, a_3 and a_7 :

$$\begin{aligned}
 a_2 = \frac{1}{2}C_8 \left(\frac{1}{12}\beta y_0^4 - \frac{1}{3}C_4 y_0^3 - C_5 y_0^2 - c_0 y_0^2 \right), \\
 a_1 = C_8, \quad a_3 = 2C_1 C_7 y_0, \\
 a_7 = -c_1(c_1 F_3(T) + F_4(T)(C_1 y_0 + C_2)). \quad (19)
 \end{aligned}$$

Then we discuss the solution of Eq. (14), the balancing procedure yields $n = 2$ for Eq. (14), therefore, we may choose

$$\begin{aligned}
 A = f(Z, T) + f_1(T)\text{sech}(W_1(T)X + W_2(T)Z \\
 + W_3(T)) + f_2(T)\text{sech}(W_1(T)X \\
 + W_2(T)Z + W_3(T))^2, \quad (20)
 \end{aligned}$$

Substituting Eq. (20) into Eq. (14), we can get $f(Z, T), f_1(T), f_2(T), W_1(T), W_2(T)$, and $W_3(T)$ by vanishing sech,

$$\begin{aligned}
 f(Z, T) &= g_1(T)Z + g_2(T), \quad f_1(T) = 0, \quad f_2(T) = \frac{12C_{10}^2 a_2^{1/3}}{a_3^{1/3}}, \\
 W_1(T) &= \frac{C_{10} a_3^{1/3}}{a_2^{1/3}}, \quad W_2(T) = \int \frac{-a_3 g_1(T) W_1(T)}{a_1} dT + C_9, \\
 W_3(T) &= \int \left(\frac{c_1^2 W_2(T)^2 F_3(T)}{C_8 W_1(T)} + \frac{c_1 W_2(T)^2 F_4(T)}{C_8 W_1(T)} C_{10} y_0 + \frac{c_1 W_2(T)^2 F_4(T) C_2}{C_8 W_1(T)} + \frac{2}{3} W_1(T)^3 C_4 y_0^3 \right. \\
 &\quad \left. - \frac{2W_1(T) C_1 C_7 y_0 g_2(T)}{C_8} + 2W_1(T)^3 c_0 y_0^2 - \frac{1}{6} W_1(T)^3 \beta y_0^4 + 2W_1(T)^3 C_5 y_0^2 \right) dT + C_{11}, \quad (21)
 \end{aligned}$$

where $g_1(T), g_2(T)$ are arbitrary functions of T , C_9, C_{10} and C_{11} are arbitrary constants. Then, we have

$$A = g_1(T)Z + g_2(T) + \frac{12C_{10}^2 a_2^{1/3}}{a_3^{1/3}} \operatorname{sech}(W_1(T)X + W_2(T)Z + W_3(T))^2. \quad (22)$$

According to Eqs. (20) and (21), we can obtain one possible approximate solution of Eq. (3) in the form

$$\begin{aligned}
 \psi &\approx U_0(y) + \epsilon U_1(y, T) + \epsilon \phi_1 + \epsilon^2 \phi_2 \\
 &= U_0(y) + \epsilon U_1(y, T) + \epsilon A G + \epsilon^2 \left(G_1 A^2 + G_2 A + G_3 \int A_Z dX \right), \quad (23)
 \end{aligned}$$

where $A, G, G_1, G_2, G_3, U_0(y)$ and $U_1(y, k)$ satisfy Eqs.(18)–(22).

When we define uncertain parameters, some new soliton solutions can be obtained as shown in Figs. 1(a) and 1(b).




Fig. 1. Evolution of solutions (23) with the parameters: $C_1 = 1, \beta = 1, y_0 = 8, \epsilon = 0.1, c_1 = 1, c_0 = 1, g_1(T) = T, C_6 = 0, C_9 = 0, g_2(T) = 0, F_1(T) = T, C_8 = 1, C_4 = 1, C_5 = 1, C_7 = 1/12, C_{11} = 0, C_{10} = 36/106^2/3, F_3(T) = 1, C_2 = 0, F_4(T) = 1/T, C_3 = 0.$

Next, we discuss the solution of Eq. (3) with the generalized cKP equation. In a similar way, equation (15) should satisfy

$$\begin{aligned}
 a_4 &= a_5 = a_6 = a_8 = a_9 = a_{10} = 0, \\
 U_{0yyy}(y)A_X G - U_{0y}(y)A_X G_{yy} - c_0 A_X G_{yy} &= 0, \\
 -G_{yy}G_y + G G_{yyy} + 2U_{0yyy}(y)G_1 - 2U_{0y}(y)G_{1yy} - 2c_0 G_{1yy} &= 0, \\
 U_{0yyy}(y)G_3 - U_{0y}(y)G_{3yy} - c_0 G_{3yy} - c_1 G_{yy} &= 0, \\
 U_{1yyy}(y, T)G - U_{1y}(y, T)G_{yy} + \beta_1 G + U_{0yyy}(y)G_2 - U_{0y}(y)G_{2yy} - c_0 G_{2yy} &= 0, \quad (24)
 \end{aligned}$$

We obtain the solution of Eq. (24) by tedious calculations,

$$\begin{aligned} G_1 &= P_5(T), \quad G_3 = c_1 P_3(T) + P_4(T)(D_1 y + D_2), \\ G &= P_5(T)P_4(T)D_1, \quad G_2 = P_3(T)(D_1 y + D_2), \\ U_0(y) &= -\frac{1}{6}\beta y^3 + \frac{1}{2}D_4 y^2 + D_5 y + D_6, \\ U_1(y, T) &= \frac{1}{2}D_1 y^2 + D_2 y + D_3 + P_1(T), \\ U_i(y, T) &= 0 \quad (i \geq 2), \end{aligned} \quad (25)$$

where $D_i, i = 1, 2, \dots, 6$ are arbitrary constants, $P_i(T), i = 1, 3, \dots, 5$ are arbitrary functions of T . Then we obtain

$$a_1 = P_5(T)P_4(T)D_1,$$

$$\begin{aligned} a_2 &= \frac{1}{24}P_5(T)P_4(T)D_1 y_0^2 (\beta y_0^2 - 4D_4 y_0 \\ &\quad - 12c_0 - 12D_5), \\ a_3 &= P_5(T)(P_4(T)P_3(T)D_1^2 + 2D_1 y_0 - 2D_2), \\ a_7 &= -c_1^2 P_3(T) - c_1 P_4(T)(D_1 y_0 + D_2), \\ a_{11} &= P_{5T}(T)P_4(T)D_1 + P_5(T)P_{4T}(T)D_1. \end{aligned} \quad (26)$$

Now we discuss the solution of Eq. (15). If we let

$$\begin{aligned} P_4(T) &= \frac{-2(D_1 y_0 - D_2)}{D_1(-1 + P_3(T)D_1)}, \\ D_4 &= \frac{1}{4} \frac{(-24 - 12c_0 y_0^2 + \beta y_0^4 - 12D_5 y_0^2)}{y_0^3}, \end{aligned} \quad (27)$$

and $P_3(T), P_5(T)$ satisfy

$$\begin{aligned} \frac{1}{2} \frac{c_1(-c_1 P_3(T)D_1 + c_1 P_3(T)^2 D_1^2 - 2D_1^2 y_0^2 + 2D_2^2)}{D_1 P_5(T)(D_1 y_0 - D_2)} &= \frac{3\sigma_1^2}{T^2}, \\ \frac{(-P_{5T}(T) + P_{5T}(T)P_3(T)D_1 - P_5(T)P_{3T}(T)D_1)}{(-1 + P_3(T)D_1)P_5(T)} &= \frac{1}{2T}, \end{aligned} \quad (28)$$

where $\sigma_1^2 = \pm 1$, which also arises in water waves. The cKP equation is known to be completely integrable.^[22] According to Eqs. (27) and (28), equation (15) becomes

$$\begin{aligned} A_T + A_{XXX} + AA_X \\ + \frac{3\sigma_1^2}{T^2} \int A_{ZZ} dX + \frac{1}{2T} A = 0, \end{aligned} \quad (29)$$

we know the solution^[16] of Eq. (49):

$$\begin{aligned} A &= 3\lambda^2 \text{sech}^2(\xi), \\ \xi &= \frac{1}{2} \left[\lambda X - \frac{\lambda \sigma_2 T Z^2}{12} \right. \\ &\quad \left. - \frac{T(\lambda^4 - \lambda \delta Z + 3\sigma_2 \delta^2)}{\lambda} + \gamma \right], \end{aligned} \quad (30)$$

where $\sigma_2 = \pm 1, \lambda, \delta, \gamma$ are integral constants. Then we obtain an approximate solution of Eq. (3):

$$\begin{aligned} \psi &\approx U_0(y) + \epsilon U_1(y, T) + \epsilon \phi_1 + \epsilon^2 \phi_2 \\ &= U_0(y) + \epsilon U_1(y, T) + \epsilon AG \\ &\quad + \epsilon^2 \left(G_1 A^2 + G_2 A + G_3 \int A_Z dX \right), \end{aligned} \quad (31)$$

where $U_0(y), U_1(y, T), G, G_1, G_2, G_3$ and A satisfy Eqs. (25)–(28) and (30).

When we define uncertain parameters, some new soliton solutions can be obtained. The evolution of solution (31) with some different parameters are shown in Figs. 2(a) and 2(b).




Fig. 2. Evolution of solution (31) with the parameters: $P_1(T) = 0, \lambda = 1, \epsilon_1 = -1, \epsilon = 0.1, \delta = 1, \gamma = 1, D_3 = 0, c_0 = 1, D_2 = 0, y_0 = 1, \beta = 0, D_1 = 1, D_6 = 0, D_5 = 0, D_4 = -0.1$.

4. The solution of BPV equation from generalized mKP equation by symmetry group method

As is well known, the generalized mKP equation is a universal model for the propagation of weakly non-linear dispersive long waves which are essentially one directional with weak transverse effects. There are many difficulties in obtaining the solution of the generalized mKP equation, but it is easier to find the solution of the mKP equation than that of the generalized mKP equation. So we give a theorem to change the generalized mKP equation into the mKP equation by a generalized symmetry group method first, then we can find a more preferable research BPV equation with the help of the theorem.

First, let

$$A(X, Z, T) = v_X, \tag{32}$$

where $v \equiv v(X, Z, T)$. Substituting Eq. (32) into the

generalized mKP equation, we have

$$a_1 v_{XT} + v_{XXXX} + a_5 v_{XX} v_X^2 + a_6 v_{XX} v_Z + a_7 v_{ZZ} = 0. \tag{33}$$

Secondly, let

$$v = \alpha + \gamma V(\xi, \eta, \tau) = \alpha + \gamma V, \tag{34}$$

where $\alpha, \gamma, \xi, \eta,$ and τ are functions of $\{X, Z, T\}$, V satisfies the following equation

$$V_{\xi\xi\xi\xi} + d_1 V_{\xi\tau} + d_3 V_{\xi\xi} V_{\xi}^2 + d_4 V_{\xi\xi} V_{\eta} + d_5 V_{\eta\eta} = 0, \tag{35}$$

where d_1, d_3, d_4 and d_5 are non-zero arbitrary constants. Substituting Eq. (34) into Eq. (33), then eliminating $V_{\xi\xi\xi\xi}$ by using Eq. (35), from that, the remained determining equations of the functions $\xi, \eta, \tau, \alpha, \gamma$ can be obtained by vanishing the coefficients of V and its derivatives. Then the general solution of the determining equations are found out by tedious calculations. The result reads

$$\begin{aligned} \xi &= c_2 \left(\frac{a_1 \tau_{0T}}{d_1} \right)^{1/3} X + \xi_0, \quad \tau = \tau_0, \quad \gamma = \delta \sqrt{\frac{d_3}{a_5}}, \quad \eta = \sqrt{\frac{a_5}{d_3}} c_2^2 \left(\frac{a_1 \tau_{0T}}{d_1} \right)^{2/3} \frac{d_4 Z}{a_6 \delta} + \eta_0, \\ \xi_0 &= \frac{(-2a_1 a_5 a_{1T} \tau_{0T} - 2a_1^2 a_5 \tau_{0TT} + 3a_1^2 a_5 T \tau_{0T})}{d_5 d_3 a_6^2 a_1^{2/3} \tau_{0T}^{2/3} d_1^{1/3}} \frac{1}{12} c_2 d_4^2 Z^2 + \xi_1 Z + \xi_2, \\ \alpha &= -\frac{1}{4} \int a_1^{1/3} \tau_{0T}^{1/3} \left(\frac{a_6^2 \xi_{0Z}^2 d_4^2 d_1^{1/3} + 4c_2 \xi_{0T} a_1^{4/3} \tau_{0T}^{1/3} a_5 d_4^2}{\tau_{0T} a_5 d_4^2 a_1 a_6} + \frac{4a_6^2 \xi_{0Z}^2 d_3 d_5 d_1^{1/3}}{\tau_{0T} a_5 d_4^2 a_1 a_6} \right) d_1^{1/3} c_2 dZ \\ &\quad - \frac{1}{2} \frac{a_6 \xi_{0Z} d_1^{1/3} X}{c_2 a_1^{1/3} \tau_{0T}^{1/3} a_5} + \alpha_0, \end{aligned} \tag{36}$$

where $\tau_0 \equiv \tau_0(T), \xi_0 \equiv \xi_0(Z, T), \xi_1 \equiv \xi_1(T), \xi_2 \equiv \xi_2(T), \eta_0 \equiv \eta_0(T), \alpha_0 \equiv \alpha_0(T), \delta = \pm 1$, and $c_2 = 1, -\frac{1}{2} + \frac{\sqrt{3}}{2}i, -\frac{1}{2} - \frac{\sqrt{3}}{2}i$. At the same time, it should satisfy four conditions:

$$\begin{aligned} a_7 a_5 d_4^2 - d_5 d_3 a_6^2 &= 0, \\ 6a_1 \eta_{0T} \sqrt{d_3 a_5 a_6^2} \delta (a_1 \tau_{0T})^{1/3} d_1^{2/3} d_4 + (-6a_1^2 a_5 \tau_{0T} d_4^2 a_6 T + 4a_1 a_5 d_4^2 a_6 a_{1T} \tau_{0T} + 4a_1^2 a_5 d_4^2 a_6 \tau_{0TT} \\ &\quad + 3a_1^2 \tau_{0T} d_4^2 a_5 T a_6) c_2^2 Z + 12d_5 d_3 a_6^3 \xi_{0Z} c_2 a_1^{2/3} \tau_{0T}^{2/3} d_1^{1/3} = 0, \\ 4\tau_{0T} a_{1T} a_5 d_5 d_3 - 3a_1 d_4^2 \tau_{0T} a_5 T + 2d_4^2 \tau_{0T} a_5 a_{1T} + 2a_1 d_4^2 a_5 \tau_{0TT} + 4a_1 \tau_{0TT} a_5 d_5 d_3 &= 0, \\ 3a_6^3 \xi_{0Z} d_4^2 d_5 d_3 a_1^{2/3} \tau_{0T}^{2/3} d_1^{1/3} \xi_{0ZZ} + 12d_5^2 d_3^2 a_6^3 \xi_{0Z} a_1^{2/3} \tau_{0T}^{2/3} d_1^{1/3} \xi_{0ZZ} + 3\tau_{0T} d_4^4 a_1^2 c_2 a_6 T \xi_{0Z} a_5 \\ &\quad + 3\tau_{0T} d_4^4 a_1^2 c_2 a_6 \xi_{0ZT} a_5 - \tau_{0T} d_4^4 a_1 c_2 a_6 \xi_{0Z} a_5 a_{1T} - d_4^4 a_1^2 c_2 a_6 \xi_{0Z} a_5 \tau_{0TT} \\ &\quad - 3\tau_{0T} d_4^4 a_1^2 c_2 a_6 \xi_{0Z} a_5 T + 6c_2 \tau_{0T} \xi_{0ZT} a_1^2 a_5 d_4^2 d_5 d_3 a_6 = 0. \end{aligned} \tag{37}$$

In summary, we obtain the following theorem.

Theorem If $U \equiv U(X, Z, T)$ is a solution of the mKP equation (35), then

$$u = \alpha + \gamma U(\xi, \eta, \tau)$$

is the solution of the generalized mKP equation (33) on the conditions of Eq. (37), where $\alpha, \gamma, \xi, \eta, \tau, \delta$ and c_2 are given by Eq. (36).

According to the Theorem, we will research the BPV equation easier with the help of the mKP and generalized mKP equation.

Next, we will write down some types of solutions of Eq. (3) with the help of the known solutions for the (2+1)-dimensional mKP equation.

As known to all, mKP equation is

$$-u_T + u_{XXX} - 6u_X u^2 - 6u_X \int u_Z dX + 3 \int u_{ZZ} dX = 0, \quad (38)$$

where $u \equiv u(X, Z, T)$.

Let

$$d_1 = -1, \quad d_3 = -6, \quad d_4 = -6, \quad d_5 = 3, \quad (39)$$

and replace $V(X, Z, T)$ with $V(\xi, \eta, \tau)$, equation (35)

becomes

$$V_{XXXX} - V_{XT} - 6V_{XX}V_X^2 - 6V_{XX}V_Z + 3V_{ZZ} = 0, \quad (40)$$

equation (40) is equivalent to Eq. (38) under the transformation $u = V_X$. So we discuss Eq. (40) firstly in the following.

We know that the mKP equation possesses the following solution:

$$u = b(1 - \tanh(b(X - 2bZ - 16b^2T))), \quad (41)$$

where b is a non-zero arbitrary constant. According to the transformation $u = V_X$ and Eq. (41), we obtain the solution of Eq. (40):

$$V = bX + \ln(\operatorname{sech}(b(-X + 2bZ + 16b^2T))), \quad (42)$$

according to the Theorem and Eq. (32), we obtain the solution of the generalized mKP equation,

$$A = \frac{1}{6a_6 a_1 \tau_0 c_2 a_5} \{6\sqrt{6}\delta\sqrt{abc_2^2 a_1 \tau_0 (-a_1 \tau_0)}^{1/3} a_6 [\tanh(b(-X + 2bZ + 16b^2T)) + 1] - 2c_2 a_1 a_5 Z (a_{1T} \tau_0 + a_1 \tau_{0TT}) - 3\xi_1 a_6^2 (-a_1 \tau_0)^{2/3}\}, \quad (43)$$

where a_5 is a non-zero arbitrary constant. In order to obtain the solution of Eq. (3), we need to know $U_0(y), U_1(y, T), G, G_1, G_2$ and G_3 , which should satisfy $a_3 = 0, a_4 = 0, a_8 = 0, a_9 = 0, a_{10} = 0, a_{11} = 0$, Eqs. (10), (12) and (37). After tedious calculations, we obtain:

$$\begin{aligned} G &= K_1 y + K_2 + Q_5(T), \quad G_1 = Q_6(T), \\ G_2 &= Q_2(T)K_6, \quad G_3 = c_1 Q_2(T) + Q_4(T)K_6, \\ U_0(y) &= -\frac{1}{6}\beta y^3 + \frac{1}{2}K_3 y^2 + K_4 y + K_5, \\ U_1(y, T) &= K_6 y + K_7 + Q_1(T), \\ U_i(y, T) &= 0 \quad (i \geq 2), \end{aligned} \quad (44)$$

where $Q_i(T), i = 1, 2, 4, 5, 6$ are arbitrary function of $T, K_i, i = 1, 2, \dots, 7$ are arbitrary constants. Then we get

$$\begin{aligned} a_2 &= \frac{1}{40}\beta K_1 y_0^5 + \frac{1}{24}y_0^4(\beta K_2 - 2K_3 K_1) \\ &\quad - \frac{1}{6}y_0^3(c_0 K_1 + K_4 K_1 + K_3 K_2) \\ &\quad - \frac{1}{2}K_2 y_0^2(c_0 + K_4), \\ a_1 &= K_1 y_0 + K_2, \quad a_5 = 4K_1^2, \end{aligned}$$

$$\begin{aligned} a_6 &= -2K_1(c_1 Q_2(T) + Q_4(T)K_6), \\ a_7 &= -c_1(c_1 Q_2(T) + Q_4(T)K_6). \end{aligned} \quad (45)$$

From that we obtain an approximate solution of Eq. (3)

$$\begin{aligned} \psi &\approx U_0(y) + \epsilon U_1(y, T) + \epsilon \phi_1 + \epsilon^2 \phi_2 \\ &= U_0(y) + \epsilon U_1(y, T) + \epsilon A G \\ &\quad + \epsilon^2 \left(G_1 A^2 + G_2 A + G_3 \int A_Z dX \right), \end{aligned} \quad (46)$$

where $A, G, G_1, G_2, G_3, U_0(y)$ and $U_1(y, k)$ satisfy Eqs. (37), (43)–(44).

5. Summary and discussion

In summary, by the reductive perturbation method the generalized (2+1)-dimensional modified Kadomtsev–Petviashvili (mKP), Kadomtsev–Petviashvili (KP) and cylindrical Kadomtsev–Petviashvili (cKP) equations are derived from the baroclinic potential vorticity (BPV) equation. Some solutions of the KP, mKP and cKP equations are derived by a direct method and symmetry group direct method. Thus the solution of the generalized mKP

equation can be obtained from the mKP equation. Furthermore, some approximate solutions of the BPV

equation are obtained from three types of generalized KP equations.

References

- [1] Chen Y and Fan E 2007 *Chin. Phys.* **16** 6
- [2] Zhang S L, Lou S Y and Qu C Z 2006 *Chin. Phys.* **15** 2765
- [3] Yang P, Chen Y and Li Z B 2008 *Chin. Phys. B* **17** 3953
- [4] Wang J and Li B 2009 *Chin. Phys. B* **18** 2109
- [5] Li J H and Lou S Y 2008 *Chin. Phys. B* **17** 747
- [6] Yao R X, Jiao X Y and Lou S Y 2009 *Chin. Phys. B* **18** 1821
- [7] Qian S P and Tian L X 2007 *Chin. Phys.* **16** 303
- [8] Zha Q L and Li Z B 2008 *Chin. Phys. B* **17** 2333
- [9] Zhang S Q, Xu G Q and Li Z B 2002 *Chin. Phys.* **11** 933
- [10] Xia T C, Zhang H Q and Yan Z Y 2001 *Chin. Phys.* **10** 694
- [11] Pedlosky J 1979 *Geophysical Fluid Dynamics* (New York: Springer)
- [12] Lou S Y, Tong B, Hu H C and Tang X Y 2006 *J. Phys. A: Math. Gen.* **39** 513
- [13] Tang X Y, Huang F and Lou S Y 2006 *Chin. Phys. Lett.* **24** 887
- [14] Gao Y and Tang X Y 2007 *Commun. Theor. Phys.* (Beijing, China) **48** 961
- [15] Li B, Chen Y and Zhang H Q 2005 *Acta Mech.* **174** 77
- [16] Tang X Y and Lou S Y 2002 *Chin. Phys. Lett.* **19** 1
- [17] Abdul-Majid Wazwaz 2008 *Appl. Math. Comput.* **204** 227
- [18] Lou S Y and Ma H C 2005 *J. Phys. A* **38** L129
- [19] Lou S Y and Ma H C 2006 *Chaos, Solitons and Fractals* **30** 804
- [20] Lou S Y, Jia M, Tang X Y and Huang F 2007 *Phys. Rev. E* **75** 056318
- [21] Lou S Y and Tang X Y 2004 *J. Math. Phys.* **45** 1020
- [22] Hereman W and Zhuang W 1995 *Acta Appl. Math.* **39** 361